

AMENDMENTS TO THE CLAIMS

1-61. (canceled)

62. (new) A method of treating premature ejaculation in a male comprising administering to the male an antidepressant via a route selected from the group consisting of:

i) mucosal administration;

ii) administration to the lungs

iii) local administration to at least a part of the male genitalia and

iv) a combination of two or more of mucosal administration, administration to the lungs and local administration to at least a part of the male genitalia.

63. (new) A method according to claim 1 wherein the antidepressant is administered prior to or during sexual activity.

64. (new) The method according to claim 1 wherein the mucosal administration is by nasal administration, buccal administration, rectal administration or combinations thereof.

65. (new) The method according to claim 1 wherein administration to the lungs is by way of inhalation.

66. (new) The method according to claim 1 wherein local administration is directly to the penis.

67. (new) A method according claim 1 wherein the antidepressant is administered by a combination of mucosal administration and local administration.

68. (new) The method according to claim 1 wherein local administration is topical administration to the penis and administration to the lungs is nasal administration.

69. (new) The method according to claim 1 wherein the antidepressant is administered in an amount of between 0.1 and 1000 mg per dose.

70. (new) The method according to claim 1 wherein a single antidepressant or single combination of antidepressants is administered by two or more routes selected from the mucosal administration, administration to the lungs and local administration to at least a part of the male genitalia.

71. (new) The method according to claim 1 wherein two or more different antidepressants or two or more different combinations of antidepressants are administered by two or more routes selected from the mucosal administration, administration to the lungs and local administration to at least a part of the male genitalia.

72. (new) The method according to claim 10 wherein the antidepressant combination includes serotonin reuptake inhibitor (SSRIs) and a monoamine oxidase inhibitor.

73. (new) The method according to claim 1 wherein the antidepressant is massaged into the skin of the male genitalia prior to intercourse for about one minute.

74. (new) The method according to claim 12 wherein the antidepressant is massaged into the skin of the male genitalia about 30 to 60 minutes prior to intercourse.

75. (new) The method according to claim 1 wherein the antidepressant is taken nasally within 30 minutes prior to intercourse.

76. (new) A composition for the treatment of premature ejaculation, said composition comprising an antidepressant formulated for mucosal administration, administration to the lungs and/or local administration to at least part of the male genitalia.

77. (new) The composition according to claim 15 wherein said antidepressant is selected from serotonin reuptake inhibitors, serotonin norepinephrine reuptake inhibitors, monoamine oxidase inhibitors, norepinephrine dopamine reuptake inhibitors, serotonin antagonist and reuptake inhibitors, norepinephrine antagonist/serotonin antagonists, synthetically derived phenylpiperazine antidepressants, antagonists of central L2- α 2 auto and heteroadrenoceptors, and mixtures thereof.

78. (new) The composition according to claim 15 wherein said antidepressant is a mixture of a serotonin reuptake inhibitor and a monoamine oxidase inhibitor.

79. (new) The composition according to claim 17 wherein said serotonin norepinephrine reuptake inhibitor is a bicyclic, tricyclic or tetracyclic antidepressant.

80. (new) The composition according to claim 15 wherein said antidepressant is selected from paroxetine, fluvoxetine, sertraline, citalopram hydrobromide, fluvoxamine, sertraline, nortriptyline hydrochloride, clomipramine hydrochloride, dothiepin hydrochloride, Imipramine hydrochloride, mianserin hydrochloride, amitriptyline hydrochloride, phenelzine sulphate, tranylcypromine

sulphate, isocarboxazid, moclobemide, serotonin and/or adrenalin update
inhibitors such as venlafaxine, nefazodone hydrochloride, trazodone, bupropion,
mirtazapine, doxepin hydrochloride, trimipramine or mixtures thereof.

81. (new) The composition according to claim 15 formulated for local
administration directly to the penis.

82. (new) The composition according to claim 15 further including an
enhancer agent selected from cyclodextrin, water, a monohydric alcohol, a
polyhydric alcohol

83. (new) The composition according to claim 15 comprising:
about 20 to about 60% water;
about 80% to about 40% alcohols
1 to 10% Hydroxypropyl- β -cyclodextrin (HPBCD) and wherein antidepressant
agent and the HPBCD are typically present in the composition in a molar ratio of
about 1:0.8 to 1:1.4 respectively.

84. (new) The composition according to claim 15 wherein the
concentration of the antidepressant is between about 1 to 10% by weight of said
composition.

85. (new) A kit comprising at least two of: an antidepressant formulated
for nasal administration; an antidepressant formulated for local administration to
at least part of the male genitalia and an antidepressant formulated for
administration to the lungs.